



Austin Cole, MD

Syndesmosis & Deltoid Repair Post-Operative Protocol

Week 0-2

Postoperative posterior splint

Non-weight bearing with crutch use, no scooter use

Post-Operative Week 2-4

Goals:

- Toe-touch progressive to full weight bearing in boot by week 4
- Continue general strengthening of the LE, UE and core as indicated

Treatment:

- Gait training
- Begin ankle AROM without resistance
 - Dorsiflexion, 75% max plantar flexion (avoid full range PF)
 - Avoid passive inversion and eversion
- Begin ankle-strengthening exercises
 - Isometrics all directions - submaximal
 - No resistance bands
- Intrinsic foot strengthening: toe curls

Post-Operative Weeks 4-8

Goals:

- Weeks 4-6 full weight bearing in walking boot
 - May sleep without boot when full weightbearing
- Weeks 6-8 wean out of boot and into shoe with Velocity Ankle Brace
- Progress ankle strength, ROM and proprioceptive exercises

Treatment:

- Normal gait patterning/training
- Progress to full AROM, all directions
- 4 way resisted ankle
- Add LE closed chain exercises, single plane
- Evaluate LE biomechanics, flexibility and strength bilaterally
- Introduce proprioceptive exercises
 - Even ground only, no BAPS, BOSU or Wobble board

Postoperative Weeks 8-12

Goals:

- Introduce step up/down progression
- Full functional ROM of the ankle
- Patient demonstrates fair static and dynamic neuromuscular control

Treatment:

- Progress closed chain exercises
- Progress proprioceptive training o Static and dynamic balance progressing to varied surfaces as patient is able

Post-Operative Weeks 12-16

Transition from Velocity Ankle Brace to traditional lace up ankle brace

Goals:

- Patient to demonstrate full strength throughout full ROM
- Patient to demonstrate fully restored static and dynamic neuromuscular control and kinesthetic awareness

Treatment:

- Begin high impact activity
- Progress to plyometric program to include dynamic, multiplanar exercises
- Increase intensity and resistance in closed chain activities
- Sport-specific drills
- Multiplanar movements

Complete functional test before 4 month follow up visit with physician.