

## Austin Cole, MD

# **Syndesmosis ORIF**

# **Post-Operative Protocol**

#### Week 0-2

Postoperative posterior splint Non-weight bearing with crutch use, no scooter use

## **Post-Operative Week 2-4**

#### Goals:

- Toe-touch progressive to full weight bearing in boot
- Continue general strengthening of the LE, UE and core as indicated

#### Treatment:

- Gait training
- Begin ankle AROM without resistance
  - o Dorsiflexion
  - o Avoid full-range plantar flexion (75% max ROM)
- Begin ankle-strengthening exercises
  - o Isometrics all directions submaximal
  - o No resistance bands
- Intrinsic foot strengthening: toe curls

## **Post-Operative Weeks 4-8**

### Goals:

- Weeks 4-6 full weight bearing in walking boot
  o May sleep out of boot when full weightbearing
- Weeks 6-8 wean out of boot and into shoe
- Progress ankle strength, ROM and proprioceptive exercises

## Treatment:

- Normal gait patterning/training
- Progress to full AROM, all directions
- Add LE closed chain exercises, single plane
- Evaluate LE biomechanics, flexibility and strength bilaterally
- Introduce proprioceptive exercises
  - o Even ground only, no BAPS, BOSU or Wobble board

#### **Postoperative Weeks 8-12**

#### Goals:

- Normalized gait
- Introduce step up/down progression
- Full functional ROM of the ankle
- Patient demonstrates fair static and dynamic neuromuscular control

## Treatment:

- Progress closed chain exercises
- Progress proprioceptive training
  - o Static and dynamic balance progressing to varied surfaces as patient is able

## Post-Operative Weeks 12-16

#### Goals:

- Patient to demonstrate full strength throughout full ROM
- Patient to demonstrate fully restored static and dynamic neuromuscular control and

### kinesthetic awareness

## Treatment:

- Begin high impact activity in lace up ankle brace
- Progress to plyometric program to include dynamic, multiplanar exercises
- Increase intensity and resistance in closed chain activities
- Sport-specific drills
- Multiplanar movements
- Progress out of ankle Lace up brace week 16

Complete functional test before four month follow up visit with physician.